

This information was sourced from www.pfdn.org

How is incontinence evaluated?

The first step toward relief is to see a doctor who has experience treating incontinence to learn what type you have. A urologist specializes in the urinary tract, and some urologists further specialize in the female urinary tract. Gynecologists and obstetricians specialize in the female reproductive tract and childbirth. A urogynecologist focuses on urinary and associated pelvic problems in women. Family practitioners and internists see patients for all kinds of health conditions. Any of these doctors may be able to help you. In addition, some nurses and other health care providers often provide rehabilitation services and teach behavioral therapies such as fluid management and pelvic floor strengthening.

To diagnose the problem, your doctor will first ask about symptoms and medical history. Your pattern of voiding and urine leakage may suggest the type of incontinence you have. Thus, many specialists begin with having you fill out a bladder diary over several days. These diaries can reveal obvious factors that can help define the problem—including straining and discomfort, fluid intake, use of drugs, recent surgery, and illness. Often you can begin treatment at the first medical visit.

Your doctor may instruct you to keep a diary for a day or more—sometimes up to a week—to record when you void. This diary should note the times you urinate and the amounts of urine you produce. To measure your urine, you can use a special pan that fits over the toilet rim. You can also use the bladder diary to record your fluid intake, episodes of urine leakage, and estimated amounts of leakage.

If your diary and medical history do not define the problem, they will at least suggest which tests you need.

Your doctor will physically examine you for signs of medical conditions causing incontinence, including treatable blockages from bowel or pelvic growths. In addition, weakness of the pelvic floor leading to incontinence may cause a condition called prolapse, where the vagina or bladder begins to protrude out of your body. This condition is also important to diagnose at the time of an evaluation.

Your doctor may measure your bladder capacity. The doctor may also measure the residual urine for evidence of poorly functioning bladder muscles. To do this, you will urinate into a measuring pan, after which the nurse or doctor will measure any urine remaining in the bladder. Your doctor may also recommend other tests:

- **Bladder stress test**—You cough vigorously as the doctor watches for loss of urine from the urinary opening.
- **Urinalysis and urine culture**—Laboratory technicians test your urine for evidence of infection, urinary stones, or other contributing causes.
- **Ultrasound**—This test uses sound waves to create an image of the kidneys, ureters, bladder, and urethra.
- **Cystoscopy**—The doctor inserts a thin tube with a tiny camera in the urethra to see inside the urethra and bladder.

- **Urodynamics**—Various techniques measure pressure in the bladder and the flow of urine.

How is incontinence treated?

Behavioral Remedies: Bladder Retraining and Kegel Exercises

By looking at your bladder diary, the doctor may see a pattern and suggest making it a point to use the bathroom at regular timed intervals, a habit called timed voiding. As you gain control, you can extend the time between scheduled trips to the bathroom. Behavioral treatment also includes Kegel exercises to strengthen the muscles that help hold in urine.

How do you do Kegel exercises?

The first step is to find the right muscles. One way to find them is to imagine that you are sitting on a marble and want to pick up the marble with your vagina. Imagine sucking or drawing the marble into your vagina.

Try not to squeeze other muscles at the same time. Be careful not to tighten your stomach, legs, or buttocks. Squeezing the wrong muscles can put more pressure on your bladder control muscles. Just squeeze the pelvic muscles. Don't hold your breath. Do not practice while urinating.

Repeat, but don't overdo it. At first, find a quiet spot to practice—your bathroom or bedroom—so you can concentrate. Pull in the pelvic muscles and hold for a count of three. Then relax for a count of three. Work up to three sets of 10 repeats. Start doing your pelvic muscle exercises lying down. This is the easiest position to do them in because the muscles do not need to work against gravity. When your muscles get stronger, do your exercises sitting or standing. Working against gravity is like adding more weight.

Be patient. Don't give up. It takes just 5 minutes a day. You may not feel your bladder control improve for 3 to 6 weeks. Still, most people do notice an improvement after a few weeks.

Some people with nerve damage cannot tell whether they are doing Kegel exercises correctly. If you are not sure, ask your doctor or nurse to examine you while you try to do them. If it turns out that you are not squeezing the right muscles, you may still be able to learn proper Kegel exercises by doing special training with biofeedback, electrical stimulation, or both.




Figure 3. Front view of bladder. Weak pelvic muscles allow urine leakage (left). Strong pelvic muscles keep the urethra closed (right).

Medicines for Overactive Bladder

If you have an overactive bladder, your doctor may prescribe a medicine to block the nerve signals that cause frequent urination and urgency.

Several medicines from a class of drugs called anticholinergics can help relax bladder muscles and prevent bladder spasms. Their most common side effect is dry mouth, although larger doses may cause blurred vision, constipation, a faster heartbeat, and flushing. Other side effects include drowsiness, confusion, or memory loss. If you have glaucoma, ask your ophthalmologist if these drugs are safe for you.

Some medicines can affect the nerves and muscles of the urinary tract in different ways. Pills to treat swelling (edema) or high blood pressure may increase your urine output and contribute to bladder control problems. Talk with your doctor; you may find that taking an alternative to a medicine you already take may solve the problem without adding another prescription.

Scientists are studying other drugs and injections that have not yet received U.S. Food and Drug Administration (FDA) approval for incontinence to see if they are effective treatments for people who were unsuccessful with behavioral therapy or pills.

Biofeedback

Biofeedback uses measuring devices to help you become aware of your body's functioning. By using electronic devices or diaries to track when your bladder and urethral muscles contract, you can gain control over these muscles. Biofeedback can supplement pelvic muscle exercises and electrical stimulation to relieve stress and urge incontinence.

Neuromodulation

For urge incontinence not responding to behavioral treatments or drugs, stimulation of nerves to the bladder leaving the spine can be effective in some patients.

Neuromodulation is the name of this therapy. The FDA has approved a device called InterStim for this purpose. Your doctor will need to test to determine if this device would be helpful to you. The doctor applies an external stimulator to determine if neuromodulation works in you. If you have a 50 percent reduction in symptoms, a surgeon will implant the device. Although neuromodulation can be effective, it is not

for everyone. The therapy is expensive, involving surgery with possible surgical revisions and replacement.

Vaginal Devices for Stress Incontinence

One of the reasons for stress incontinence may be weak pelvic muscles, the muscles that hold the bladder in place and hold urine inside. A pessary is a stiff ring that a doctor or nurse inserts into the vagina, where it presses against the wall of the vagina and the nearby urethra. The pressure helps reposition the urethra, leading to less stress leakage. If you use a pessary, you should watch for possible vaginal and urinary tract infections and see your doctor regularly.

Injections for Stress Incontinence

A variety of bulking agents, such as collagen and carbon spheres, are available for injection near the urinary sphincter. The doctor injects the bulking agent into tissues around the bladder neck and urethra to make the tissues thicker and close the bladder opening to reduce stress incontinence. After using local anesthesia or sedation, a doctor can inject the material in about half an hour. Over time, the body may slowly eliminate certain bulking agents, so you will need repeat injections. Before you receive an injection, a doctor may perform a skin test to determine whether you could have an allergic reaction to the material. Scientists are testing newer agents, including your own muscle cells, to see if they are effective in treating stress incontinence. Your doctor will discuss which bulking agent may be best for you.

Surgery for Stress Incontinence

In some women, the bladder can move out of its normal position, especially following childbirth. Surgeons have developed different techniques for supporting the bladder back to its normal position. The three main types of surgery are retropubic suspension and two types of sling procedures.

Retropubic suspension uses surgical threads called sutures to support the bladder neck. The most common retropubic suspension procedure is called the Burch procedure. In this operation, the surgeon makes an incision in the abdomen a few inches below the navel and then secures the threads to strong ligaments within the pelvis to support the urethral sphincter. This common procedure is often done at the time of an abdominal procedure such as a hysterectomy.

Sling procedures are performed through a vaginal incision. The traditional sling procedure uses a strip of your own tissue called fascia to cradle the bladder neck. Some slings may consist of natural tissue or man-made material. The surgeon attaches both ends of the sling to the pubic bone or ties them in front of the abdomen just above the pubic bone.

Midurethral slings are newer procedures that you can have on an outpatient basis. These procedures use synthetic mesh materials that the surgeon places midway along the urethra. The two general types of midurethral slings are retropubic slings, such as the transvaginal tapes (TVT), and transobturator slings (TOT). The surgeon makes small incisions behind the pubic bone or just by the sides of the vaginal opening as

well as a small incision in the vagina. The surgeon uses specially designed needles to position a synthetic tape under the urethra. The surgeon pulls the ends of the tape through the incisions and adjusts them to provide the right amount of support to the urethra.

If you have pelvic prolapse, your surgeon may recommend an anti-incontinence procedure with a prolapse repair and possibly a hysterectomy.



Figure 4. Side view. Supporting sutures in place following retropubic or transvaginal suspension (left). Sling in place, secured to the pubic bone (center). The ends of the transobturator tape supporting the urethra are pulled through incisions in the groin to achieve the right amount of support (right). The tape ends are removed when the incisions are closed.

Recent women's health studies performed with the Urinary Incontinence Treatment Network (UITN) compared the suspension and sling procedures and found that, 2 years after surgery, about two-thirds of women with a sling and about half of women with a suspension were cured of stress incontinence. Women with a sling, however, had more urinary tract infections, voiding problems, and urge incontinence than women with a suspension. Overall, 86 percent of women with a sling and 78 percent of women with a suspension said they were satisfied with their results.

Talk with your doctor about whether surgery will help your condition and what type of surgery is best for you. The procedure you choose may depend on your own preferences or on your surgeon's experience. Ask what you should expect after the procedure. You may also wish to talk with someone who has recently had the procedure. Surgeons have described more than 200 procedures for stress incontinence, so no single surgery stands out as best.

Catheterization

If you are incontinent because your bladder never empties completely—overflow incontinence—or your bladder cannot empty because of poor muscle tone, past surgery, or spinal cord injury, you might use a catheter to empty your bladder. A catheter is a tube that you can learn to insert through the urethra into the bladder to drain urine. You may use a catheter once in a while or on a constant basis, in which case the tube connects to a bag that you can attach to your leg. If you use an indwelling—long-term—catheter, you should watch for possible urinary tract infections.

Other Helpful Hints

Many women manage urinary incontinence with menstrual pads that catch slight leakage during activities such as exercising. Also, many people find they can reduce incontinence by restricting certain liquids, such as coffee, tea, and alcohol.

Finally, many women are afraid to mention their problem. They may have urinary incontinence that can improve with treatment but remain silent sufferers and resort to wearing absorbent undergarments, or diapers. This practice is unfortunate, because diapering can lead to diminished self-esteem, as well as skin irritation and sores. If you are relying on diapers to manage your incontinence, you and your family should discuss with your doctor the possible effectiveness of treatments such as timed voiding and pelvic muscle exercises.

Points to Remember

- Urinary incontinence is common in women.
- All types of urinary incontinence are treatable.
- Incontinence is treatable at all ages.
- You need not be embarrassed by incontinence.

Hope through Research

The National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) has many research programs aimed at finding treatments for urinary disorders, including urinary incontinence. The NIDDK is sponsoring the Urinary Incontinence Treatment Network (UITN), a consortium of urologists and urogynecologists who are evaluating and comparing treatment methods for stress and mixed incontinence in women. The goal of the first study, completed in 2007, was to learn which treatment methods have the best short- and long-term outcomes for treating stress urinary incontinence in women. Ongoing studies focus on treatments for urge incontinence and minimally invasive treatments for stress incontinence.

The National Institute of Child Health and Human Development also supports research in the area of pelvic health. The Pelvic Floor Disorders Network (PFDN) was formed in 2001 to do research to improve the care and daily lives of women with pelvic organ prolapse and bladder and bowel control problems.